

Name \_\_\_\_\_ Date \_\_\_\_\_ Period \_\_\_\_\_

**Worksheet 1.1—Limits & Continuity****Short Answer:** Show all work. Unless stated otherwise, no calculator permitted.

1. Explain in your own words what is meant by the equation  $\lim_{x \rightarrow 2} f(x) = 4$ . Is it possible for this statement to be true and yet  $f(2) = 5$ ? Explain. What graphical manifestation would  $f(x)$  have at  $x = 2$ ? Sketch a possible graph of  $f(x)$ .

2. Explain what it means to say that  $\lim_{x \rightarrow 1^-} f(x) = 3$  and  $\lim_{x \rightarrow 1^+} f(x) = 6$ . What graphical manifestation would  $f(x)$  have at  $x = 1$ ? Sketch a possible graph of  $f(x)$ .

3. Explain the meaning of each of the following, then sketch a possible graph of a function exhibiting the indicated behavior.

(a)  $\lim_{x \rightarrow -2} f(x) = \infty$

(b)  $\lim_{x \rightarrow -3^+} g(x) = -\infty$ .

4. For  $f(x) = \frac{x^2 + x - 20}{x^2 - 16}$ , algebraically determine the following:

(a)  $f(4)$

(b)  $\lim_{x \rightarrow 4^-} f(x)$

(c)  $\lim_{x \rightarrow 4^+} f(x)$

(d)  $\lim_{x \rightarrow 4} f(x)$

(e)  $\lim_{x \rightarrow -4} f(x)$

(f)  $\lim_{x \rightarrow 0^-} f(x)$

(g)  $\lim_{x \rightarrow 1} f(x)$

(h)  $\lim_{x \rightarrow -1} f(x)$

5. Using the definition of continuity, determine whether the graph of  $f(x) = \frac{x^2 + x}{x^3 + 2x^2 - 3x}$  is continuous at the following. Justify.

(a)  $x = 0$

(b)  $x = 1$

(c)  $x = 2$

6. For  $f(x) = \begin{cases} -x^2, & x < 0 \\ 0.001, & x = 0 \\ \sqrt{x}, & x > 0 \end{cases}$ , algebraically determine the following:

(a)  $f(0)$       (b)  $\lim_{x \rightarrow 0^-} f(x)$       (c)  $\lim_{x \rightarrow 0^+} f(x)$       (d)  $\lim_{x \rightarrow 0} f(x)$       (e) continuity of  $f$  at  $x = 0$ . Justify.

7. Evaluate each of the following continuous functions at the indicated  $x$ -value:

(a)  $\lim_{\theta \rightarrow \frac{11\pi}{6}} \sin \theta =$       (b)  $\lim_{x \rightarrow 6} 2^x =$       (c)  $\lim_{x \rightarrow 0} (57x^{85} - 2x^{45} + 100x^{11} - 99999x + 5) =$

8. Evaluate each of the following:

(a)  $\lim_{x \rightarrow \frac{\pi}{2}^-} \tan x =$

(b)  $\lim_{x \rightarrow \frac{\pi}{2}^+} \tan x =$


(c)  $\lim_{x \rightarrow \frac{\pi}{2}} \tan x =$

(d)  $\lim_{x \rightarrow -5^-} \frac{-2}{x+5} =$

(e)  $\lim_{x \rightarrow -5^+} \frac{-2}{x+5} =$

(f)  $\lim_{x \rightarrow -5} \frac{-2}{x+5} =$

9. For the function  $f$  whose graph is given at below, evaluate the following, if it exists. If it does not exist, explain why.



(a)  $\lim_{x \rightarrow 3} f(x) =$

(b)  $\lim_{x \rightarrow 7} f(x) =$

(c)  $\lim_{x \rightarrow -4} f(x) =$


(d)  $\lim_{x \rightarrow -9^-} f(x) =$

(e)  $\lim_{x \rightarrow -9^+} f(x) =$

(f)  $\lim_{x \rightarrow -9} f(x) =$

(g) What are the equations of the vertical asymptotes?

10. A patient receives a 150-mg injection of a drug every four hours. The graph at right shows the amount  $C(t)$  of the drug in the bloodstream after  $t$  hours. Approximate  $\lim_{t \rightarrow 12^-} C(t)$  and  $\lim_{t \rightarrow 12^+} C(t)$ , then explain in a complete sentence the significance/meaning of these one-sided limits in terms of the injections at  $t = 12$  hours.



11. **(Calculator Permitted)** Sketch the graph of the function  $f(x) = \frac{1}{1 + 2^{1/x}}$  in the space below, then evaluate each, if it exists. If it does not exist, explain why. Name the type and location of any discontinuity.

$$(a) \lim_{x \rightarrow 0^-} f(x) =$$

$$(b) \lim_{x \rightarrow 0^+} f(x) =$$

$$(c) \lim_{x \rightarrow 0} f(x) =$$

$$(d) f(0) =$$

12. Using the definition of continuity at a point, discuss the continuity of the following function. Justify.

$$f(x) = \begin{cases} 2 - x, & x < -1 \\ x, & -1 \leq x < 1 \\ (x - 1)^2, & x \geq 1 \end{cases}$$

13. For  $f(x) = \begin{cases} 3ax - b, & x < 1 \\ 5, & x = 1 \\ 2a\sqrt{x} + b, & x > 1 \end{cases}$ , find the values of  $a$  and  $b$  such that  $f(x)$  is continuous at  $x = 1$ . Show the work that leads to your answer.

14. **(Calculator permitted)** Fill in the table for the following function, then use the numerical evidence (to 3 decimal places) to evaluate the indicated limit. (Be sure you're in radian mode)

$$f(x) = \frac{\sin(3x)}{x}$$

$x$	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$							




Based on the numeric evidence above,  $\lim_{x \rightarrow 0} f(x) =$

**Multiple Choice:** Put the Capital Letter of the correct answer in the blank to the left of the number. Be sure to show any work/analysis.

\_\_\_\_\_ 15.  $\lim_{x \rightarrow 0^-} \left(1 - \frac{1}{x}\right) =$   
 (A) 1      (B) 2      (C)  $-\infty$       (D) 0      (E)  $\infty$


\_\_\_\_\_ 16. Find  $\lim_{x \rightarrow 1} f(x)$  if  $f(x) = \begin{cases} 3-x, & x \neq 1 \\ 1, & x = 1 \end{cases}$   
 (A) 2      (B) 1      (C)  $\frac{3}{2}$       (D) 0      (E) DNE

\_\_\_\_\_ 17. For which of the following does  $\lim_{x \rightarrow 4} f(x)$  exist?



(A) I only    (B) II only    (C) III only    (D) I and II only    (E) I and III only

\_\_\_\_\_ 18. If  $f(x) = \begin{cases} \ln x, & 0 < x \leq 2 \\ x^2 \ln 2, & 2 < x \leq 4 \end{cases}$ , then  $\lim_{x \rightarrow 2} f(x)$  is  
 (A)  $\ln 2$       (B)  $\ln 8$       (C)  $\ln 16$       (D) 4      (E) nonexistent



Use the graph of  $f(x)$  above to answer questions 19 - 22.

\_\_\_\_\_ 19.  $\lim_{x \rightarrow 7} f(x) =$

- (A) 1      (B) 2      (C) -1      (D) 4      (E) DNE

\_\_\_\_\_ 20.  $\lim_{x \rightarrow 0^-} f(x) =$

- (A) 1      (B) 2      (C) -1      (D) 4      (E) DNE

\_\_\_\_\_ 21.  $\lim_{x \rightarrow 2} f(x) =$

- (A) 2      (B) 3      (C) -1      (D) 4      (E) DNE

\_\_\_\_\_ 22. Which of the following regarding  $f(x)$  at  $x = 5$  true?

- I.  $\lim_{x \rightarrow 5^-} f(x) = 3$   
 II.  $\lim_{x \rightarrow 5^+} f(x) = f(5)$   
 III.  $f(x)$  is continuous at  $x = 5$

- (A) I only    (B) II only    (C) I and II only    (D) II and III only    (E) I, II, and III

\_\_\_\_\_ 23. If  $f(x) = \begin{cases} ae^x + b, & x < 0 \\ 4, & x = 0 \\ bx - 2a, & x > 0 \end{cases}$ , then the value of  $b$  that makes  $f(x)$  continuous at  $x = 0$  is

(A) 2      (B) -2      (C) 4      (D) 6      (E) no such value exists

\_\_\_\_\_ 24. If  $f(x) = \frac{1}{x-2}$  and  $\lim_{x \rightarrow (-k+1)} f(x)$  does not exist, then  $k =$

(A) 2      (B) 3      (C) 1      (D) -2      (E) -1

\_\_\_\_\_ 25. The function  $f(x) = \begin{cases} \frac{x^2}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$

- (A) is continuous for all  $x$   
(B) has a removable point discontinuity at  $x = 0$   
(C) has a non-removable oscillation discontinuity at  $x = 0$   
(D) has a non-removable infinite discontinuity at  $x = 0$   
(E) has a non-removable jump discontinuity at  $x = 0$

\_\_\_\_\_ 26. If  $f(x) = \begin{cases} \frac{x^2 - x}{2x}, & x \neq 0 \\ k, & x = 0 \end{cases}$  is continuous at  $x = 0$ , then  $k =$

(A) -1      (B)  $-\frac{1}{2}$       (C) 0      (D)  $\frac{1}{2}$       (E) 1